Plant Molecular Biology **24**: 981–988, 1994.

Contents of Volume 24

Vol. 24 No. 1 January (I) 1994

Mini-review

Physiological roles for secondary metabolites in plants: some progress, many outstanding problems 1
M.J.C. Rhodes

Research articles

Studies on wheat acetyl CoA carboxylase and the cloning of a partial cDNA 21
K.M. Elborough, J.W. Simon, R. Swinhoe, A.R. Ashton, A.R. Slabas

Molecular cloning of two different cDNAs for maize acetyl CoA carboylase 35
A.R. Ashton, C.L.D. Jenkins, P.R. Whitfeld

Production of fertile transgenic maize by electroporation of suspension culture cells 51
C.M. Laursen, R.A. Krzyzek, C.E. Flick, P.C. Anderson, T.M. Spencer

Photosystem II genes *IsiA, PsbDI* and *PsbC* in *Anabaena* Sp. PCC 7120: cloning, sequencing and the transcriptional regulation in iron-stressed and iron-replete cells 63
K. Leonhardt and N.A. Straus

Pea (*Pisum sativum* L.) seed isolectins 1 and 2 and pea root lectin result from carboxypeptidase-like processing of a single gene product 75
F.J. Hoedemaeker, M. Richardson, C.L. Díaz, B.S. de Pater, J.W. Kijne

Expression of biologically active hordothionins in tobacco. Effects of pre- and pro-sequences at the amino and carboxyl termini of the hordothionin precursor on mature protein expression and sorting 83
D.E.A. Florack, W.G. Dirkse, B. Visser, F. Heidekamp, W.J. Stiekema

The highly expressed tapetum-specific A9 gene is not required for male fertility in *Brassica napus* 97
K. Turgut, T. Barsby, M. Craze, J. Freeman, R. Hodge, W. Paul, R. Scott

A 20 nucleotide upstream element is essential for the nopaline synthase (*nos*) promoter activity 105
Y. Kim, K. Buckley, M.A. Costa, G. An

Isolation of a ubiquitin-ribosomal protein gene (*ubi 3*) from potato and expression of its promoter in transgenic plants 119
J.E. Garbarino, W.R. Belknap

Subunit 6 of the F_o-ATP synthase complex from cytoplasmic male-sterile radish: RNA editing and NH_2-terminal protein sequencing 129
S. Krishnasamy, R.A. Grant, C.A. Makaroff

Members of a broadbean nodulin family with partial homologies to the alfalfa nodulin 25 are composed of two types of amino acid repeats flanked by unique amino acid sequence termini 143
H. Küster, A.M. Perlick, A. Pühler

Characterization and sequencing of cDNA clone encoding the phloem protein PP2 of *Cucurbita pepo* 159
M-B. Wang, D. Boulter, J.A. Gatehouse

Purification and partial characterization of a glycoprotein from pea (*Pisum sativum*) with receptor activity for rhicadhesin, an attachment protein of Rhizobiaceae 171
S. Swart, T.J.J. Logman, G. Smit, B.J.J. Lugtenberg, J.W. Kijne

Expression of nitrate assimilation related genes in *Chlamydomonas reinhardtii* 185
A. Quesada, E. Fernández

Expression of one of the members of the *Arabidopsis* chaperonin 60β gene family is developmentally regulated and wound-repressible 195
E. Zabaleta, N. Assad, A. Oropeza, G. Salerno, L. Herrera-Estrella

Characterization of a weak allele of the *GL1* gene of *Arabidopsis thaliana* 203
J.J. Esch, D.G. Oppenheimer, M.D. Marks

The presequence of the precursor to the nucleus-encoded 30 kDa protein of photosystem II in *Euglena gracilis* Z includes two hydrophobic domains 209
Y. Shigemori, J. Inagaki, H. Mori, M. Nishimura, S. Takahashi, Y. Yamamoto

Functional loss of all *ndh* genes in an otherwise relatively unaltered plastid genome of the holoparasitic flowering plant *Cuscuta reflexa* 217
G. Haberhausen, K. Zetsche

Characterization of a mRNA that accumulates during development of oilseed rape pods 223
S.A. Coupe, J.E. Taylor, P.G. Isaac, J.A. Roberts

Short communications

The heat shock response of an antarctic alga is evident at 5 °C 229
M.E. Vayda, M-L. Yuan

Two ABA-responsive proteins from pea (*Pisum sativum* L.) are closely related to intracellular pathogenesis-related proteins 235
E.A. Iturriaga, M.J. Leech, D.H.P. Baratt, T.L. Wang

Expression of TGMV antisense RNA in transgenic tobacco inhibits replication of BCTV but not ACMV geminiviruses 241
E.R. Bejarano, C.P. Lichtenstein

Structure and expression of a cDNA encoding an RNA helicase-like protein in tobacco 249
H. Itadani, M. Sugita, M. Sugiura

The gene for ribosomal protein L27 is located on the plastid rather than the nuclear genome of the chlorophyll *c*-containing alga *Pleurochrysis carterae* 253
S. Fujiwara, M. Kawachi, I. Inouye, J. Someya

Sequence announcement 259

Volume 24 No. 2 January (II) 1994

Research articles

Tissue-specific and temporal regulation of a β-conglycinin gene: roles of the RY repeat and other *cis*-acting elements 261
T. Fujiwara, R.N. Beachy

Palindromic repeated sequences (PRSs) in the mitochondrial genome of rice: evidence for their insertion after divergence of the genus *Oryza* from the other Gramineae 273
M. Nakazono, A. Kanno, N. Tsutsumi, A. Hirai

Methotrexate does not block import of a DHFR fusion protein into chloroplasts 283
T. America, J. Hageman, A. Guéra, F. Rook, K. Archer, K. Keegstra, P. Weisbeek

Characterization of four new β-tubulin genes and their expression during male flower development in maize (*Zea mays* L.) 295
R. Villemur, N.A. Haas, C.M. Joyce, D.P. Snustad, C.D. Silflow

Fertile transgenic barley by particle bombardment of immature embryos 317
A. Ritala, K. Aspegren, U. Kurtén, M. Salmenkallio-Marttila, L. Mannonen, R. Hannus, V. Kauppinen, T.H. Teeri, T.-M. Enari

A seed-specific *Brassica napus* oleosin promoter interacts with a G-box-specific protein and may be bi-directional 327
J.S. Keddie, M. Tsiantis, P. Piffanelli, R. Cella, P. Hatzopoulos, D.J. Murphy

Expression of mouse metallothionein-I gene confers cadmium resistance in transgenic tobacco plants 341
A. Pan, M. Yang, F. Tie, L. Li, Z. Chen, B. Ru

Primary structure and expression of mRNAs encoding basic chitinase and 1,3-β-glucanase in potato 353
L. Beerhues, E. Kombrink

A novel carotenoid biosynthesis gene coding for ζ-carotene desaturase: functional expression, sequence and phylogenetic origin 369
H. Linden, N. Misawa, T. Saito, G. Sandmann

Control of gene expression in plant cells using a 434:VP16 chimeric protein 381
R.J. Wilde, S.E. Cooke, W.J. Brammar, W. Schuch

Short communications

Structural organization of *de novo* purine biosynthesis enzymes in plants: 5-aminoimidazole ribonucleotide carboxylase and 5-aminoimidazole-4-*N*-succinocarboxamide ribonucleotide synthetase cDNAs from *Vigna aconitifolia* 389
K.A. Chapman, A.J. Delauney, J.H. Kim, D.P.S. Verma

A microsatellite marker in *Helianthus annuus?* L. 397
D. Brunel

Stable transformation and long-term expression of the *gusA* reporter gene in callus lines of perennial ryegrass (*Lolium perenne* L.) 401
H.M. van der Maas, E.R. de Jong, S. Rueb, L.A.M. Hensgens, F.A. Krens

Volume 24 No. 3 February (I) 1994

Mini-review

MAP kinases: universal multi-purpose signaling tools 407
C. Jonak, E. Heberle-Bors, H. Hirt

Research articles

Hypoxically inducible barley alanine aminotransferase: cDNA cloning and expression analysis 417
D.G. Muench, A.G. Good

Cloning and characterization of the gene for phytoene desaturase (*Pds*) from tomato *(Lycopersicon esculentum)* 429
V. Mann, I. Pecker, J. Hirschberg

Cloning and characterization of genes for tetrapyrrole biosynthesis from the cyanobacterium *Anacystis nidulans* R2 435
M.C. Jones, J.M. Jenkins, A.G. Smith, C.J. Howe

Addition of A- and U-rich sequence increases the splicing efficiency of a deleted form of maize intron 449
K.R. Luehrsen, V. Walbot

Identification of a homeobox-containing gene with enhanced expression during soybean (*Glycine max* L.) somatic embryo development 465
H. Ma, M.D. McMullen, J.J. Finer

Isolation of high-molecular-weight plant DNA for DNA damage quantitation: relative effects of solar 297 nm UVB and 365 nm radiation 475
F.E. Quaite, J.C. Sutherland, B.M. Sutherland

Characterization of a novel *cis*-acting element that is responsive to a fungal elicitor in the promoter of a tobacco class I chitinase gene 485
Y. Fukuda, H. Shinshi

***In vitro* characterization of a cassette to accumulate multiple proteins through synthesis of a self-processing polypeptide** 495
J.F. Marcos, R.N. Beachy

Selection for kanamycin resistance in transformed petunia cells leads to the co-amplification of a linked gene 505
J.D. Jones, S.C. Weller, P.B. Goldsbrough

Structural and functional analysis of an *Opaque-2*-related gene from sorghum 515
L. Pirovano, S. Lanzini, H. Hartings, N. Lazzaroni, V. Rossi, R. Joshi, R.D. Thompson, F. Salamini, M. Motto

Short communications

Cloning and characterization of a tomato GTPase-like gene related to yeast and *Arabidopsis* genes involved in vesicular transport 525
C. Davies

Complementation of a *Chlamydomonas reinhardtii* mutant using a genomic cosmid library 533
S. Purton, J.-D. Rochaix

Nucleotide sequence and expression of a ripening and water stress-related cDNA from tomato with homology to the MIP class of membrane channel proteins 539
R.G. Fray, A. Wallace, D. Grierson, G.W. Lycett

Isolation and characterization of a rice cDNA similar to the S-phase-specific *cyc07* gene 545
S.-I. Kidou, M. Umeda, T. Tsuge, A. Kato, H. Uchimiya

Volume 24 No. 4 February (II) 1994

Research articles

Structure and regulation of an ABA- and desiccation-responsive gene from the resurrection plant *Craterostigma plantagineum* 549
D. Michel, A. Furini, F. Salamini, D. Bartels

Regulation of the *rab*17 gene promoter in transgenic *Arabidopsis* wild-type, ABA-deficient and ABA-insensitive mutants 561
J. Vilardell, J.M. Martínez-Zapater, A. Goday, C. Arenas, M. Pagès

The *S11* and *S13* self incompatibility alleles in *Solanum chacoense* Bitt. are remarkably similar 571
M.K. Saba-El-Leil, S. Rivard, D. Morse, M. Cappadocia

Conserved gene clusters in the highly rearranged chloroplast genomes of *Chlamydomonas moewusii* and *Chlamydomonas reinhardtii* 585
E. Boudreau, C. Otis, M. Turmel

Cloning of two gibberellin-regulated cDNAs from *Arabidopsis thaliana* by subtractive hybridization: expression of the tonoplast water channel, γ-TIP, is increased by GA_3 603
A.L. Phillips, A.K. Huttly

Structure and light-regulated expression of the *gsa* gene encoding the chlorophyll biosynthetic enzyme, glutamate 1-semialdehyde aminotransferase, in *Chlamydomonas reinhardtii* 617
G.L. Matters, S.I. Beale

Truncated presequences of mitochondrial F_1-ATPase β subunit from *Nicotiana plumbaginifolia* transport CAT and GUS proteins into mitochondria of transgenic tobacco 631
F. Chaumont, M. de Castro Silva Filho, D. Thomas, S. Leterme, M. Boutry

Identification of conserved domains in the Δ12 desaturases of cyanobacteria 643
T. Sakamoto, H. Wada, I. Nishida, M. Ohmori, N. Murata

Homologues of wheat ubiquitin-conjugating enzymes – *Ta*UBC1 and *Ta*UBC4 are encoded by small multigene families in *Arabidopsis thaliana* 651
M.L. Sullivan, T.B. Carpenter, R.D. Vierstra

Gene isolation through genomic complementation using an indexed library of *Chlamydomonas reinhardtii* DNA 663
H. Zhang, P.L. Herman, D.P. Weeks

Short communications

Nucleotide sequence of a cDNA clone encoding the precursor of the peridinin-chlorophyll *a*-binding protein from the dinoflagellate *Symbiodimium* sp. 673
B.J. Norris, D.J. Miller

Organization and transcription of the class I phycoerythrin genes of the marine cyanobacterium *Synechococcus* sp. WH7803 679
J. Newman, N.H. Mann, N.G. Carr

Integration of CAPS markers into the RFLP map generated using recombinant inbred lines of *Arabidopsis thaliana* 685
P. Jarvis, C. Lister, V. Szabo, C. Dean

A homologue of the 65 kDa regulatory subunit of protein phosphatase 2A in early pea (*Pisum sativum* L.) embryos 689
I.M. Evans, T. Fawcett, D. Boulter, A.P. Fordham-Skelton

Sequence announcements 697

List of reviewers 699

Volume 24 No. 5 March (I) 1994

Research articles

The 5′ region of *Arabidopsis thaliana cor15a* has *cis*-acting elements that confer cold-, drought- and ABA-regulated gene expression 701
S.S. Baker, K.S. Wilhelm, M.F. Thomashow

The soybean SAUR open reading frame contains a *cis* element responsible for cycloheximide-induced mRNA accumulation 715
Y. Li, T.J. Strabala, G. Hagen, T.J. Guilfoyle

Genes encoding acidic and basic class III β-1,3- glucanases are expressed in tomato plants upon viroid infection 725
C. Domingo, V. Conejero, P. Vera

Isolation and characterization of a cDNA from flowers of *Cynara cardunculus* encoding cyprosin (an aspartic proteinase) and its use to study the organ-specific expression of cyprosin 733
M.C. Cordeiro, Z.-T. Xue, M. Pietrzak, M.S. Pais, P.E. Brodelius

Cloning and molecular analysis of structural genes involved in flavonoid and stilbene biosynthesis in grape (*Vitis vinifera* L.) 743
F. Sparvoli, C. Martin, A. Scienza, G. Gavazzi, C. Tonelli

Differential expression of two calmodulin genes in response to physical and chemical stimuli 757
J.R. Botella, R.N. Arteca

Isolation of chalcone synthase and chalcone isomerase cDNAs from alfalfa (*Medicago sativa* L.): highest transcript levels occur in young roots and root tips 767
H.I. McKhann, A.M. Hirsch

Directed disruption of the *Chlamydomonas* chloroplast *psbK* gene destabilizes the photosystem II reaction center complex 779
Y. Takahashi, H. Matsumoto, M. Goldschmidt-Clermont, J.-D. Rochaix

The presence of enhancers adjacent to the *Ac* promoter increases the abundance of transposase mRNA and alters the timing of *Ds* excision in *Arabidopsis* 789
L. Balcells, G. Coupland

Short communications

A cold-induced gene from *Medicago sativa* encodes a bimodular protein similar to developmentally regulated proteins 799
Y. Castonguay, S. Laberge, P. Nadeau, L.-P. Vézina

Sequence and functional analyses of the rice gene homologous to the maize *Vp1* 805
T. Hattori, T. Terada, S. Hamasuna

Rps3 and _rpl16_ genes do not overlap in _Oenothera_ mitochondria: GTG as a potential translation initiation codon in plant mitochondria? 811
H. Bock, A. Brennicke, W. Schuster

A cold-regulated cyanobacterial gene cluster encodes RNA-binding protein and ribosomal protein S21 819
N. Sato

Molecular cloning of mannose-binding lectins from _Clivia miniata_ 825
E.J.M. van Damme, K. Smeets, F. van Leuven, W.J. Peumans

Errata 831

Volume 24 No. 6 March (II) 1994

Research articles

Molecular analysis of the aspartate kinase-homoserine dehydrogenase gene from _Arabidopsis thaliana_ 835
M. Ghislain, V. Frankard, D. Vandenbossche, B.F. Matthews, M. Jacobs

A cereal haemoglobin gene is expressed in seed and root tissues under anaerobic conditions 853
E.R. Taylor, X.Z. Nie, A.W. MacGregor, R.D. Hill

Meristem-specific gene expression directed by the promoter of the S-phase-specific gene, _cyc07_, in transgenic _Arabidopsis_ 863
M. Ito, T. Sato, H. Fukuda, A. Komamine

Low-temperature-responsive barley genes have different control mechanisms 879
M.A. Dunn, N.J. Goddard, L. Zhang, R.S. Pearce, M.A. Hughes

The $tRNA^{Ser}$-isoacceptors and their genes in _Nicotiana rustica_: genome organization, expression _in vitro_ and sequence analyses 889
T. Teichmann, C. Urban, H. Beier

Cloning and characterization of _Tag 1_, a tobacco anther β-1,3-glucanase expressed during tetrad dissolution 903
P.A. Bucciaglia, A.G. Smith

Structural analysis of two length variants of the rDNA intergenic spacer from _Eruca sativa_ 915
M. Lakshmikumaran, M.S. Negi

Analysis of transgenic tobacco plants expressing a truncated form of a potyvirus coat protein nucleotide sequence 929
L. Silva-Rosales, J.A. Lindbo, W.G. Dougherty

Structure and expression of a cDNA encoding a histone H2A from _Euglena gracilis_ 941
A. Saint-Guily, M.-L. Schantz, R. Schantz

Molecular analysis of wild-type and mutant alleles at the _Opaque-2_ regulatory locus of maize reveals different mutations and types of _O2_ product 949
L. Bernard, P. Ciceri, A. Viotti

Short communications

Characterization and accumulation pattern of an mRNA encoding an abscission-related β-1,4-glucanase from leaflets of *Sambucus nigra* 961
J.E. Taylor, S.A. Coupe, S. Picton, J.A. Roberts

Nucleotide sequence of a rice cDNA similar to a maize NADP-dependent malic enzyme 965
T. Fushimi, M. Umeda, T. Shimazaki, A. Kato, K. Toriyama, H. Uchimiya

Sequence of a plant cDNA from *Vicia faba* encoding a novel Ran-related GTP-binding protein 969
G. Saalbach, V. Christov

Announcements 973

Erratum 975

Instructions to Authors 977

Contents of Volume 24 981

Author Index 991